

ISSN: 0256-307X

中国物理快报

Volume 28 Number 11 November 2011

A Series Journal of the Chinese Physical Society
Distributed by IOP Publishing

Online: <http://iopscience.iop.org/cpl>
<http://cpl.iphy.ac.cn>

CHINESE PHYSICAL SOCIETY
Institute of **Physics** PUBLISHING

Rogue Wave, Breathers and Bright-Dark-Rogue Solutions for the Coupled Schrödinger Equations

GUO Bo-Ling(郭柏灵)¹, LING Li-Ming(凌黎明)^{1,2**}

¹*Institute of Applied Physics and Computational Mathematics, Beijing 100088*

²*The Graduate School of China Academy of Engineering Physics, Beijing 100088*

(Received 12 May 2011)

We construct explicit rogue wave solutions, breather solitons, and rogue-bright-dark solutions for the coupled non-linear Schrödinger equations by the Darboux transformation.

PACS: 02.30.Ik, 02.30.Hq, 03.75.Nt

DOI:10.1088/0256-307X/28/11/110202

The coupled non-linear Schrödinger (CNLS/NLS) equations are a significant model in areas such as non-linear optics,^[1] bio-physics,^[2] multi-component Bose-Einstein condensates at zero temperature,^[3] finance^[4] and oceanics.^[5] However, they become integrable for a special choice of the parameter (Manakov^[6] model). On the other hand, the Manakov model has appeared in a Kerr-type approximation of photorefractive crystals. Therefore this equation can explicitly describe some physical phenomena. There is lots of the research about these kinds of system.^[7–9]

In this Letter, we consider an integrable coupled Schrödinger equation

$$iq_{1t} + q_{1xx} + 2(|q_1|^2 + |q_2|^2)q_1 = 0; \quad (1a)$$

$$iq_{2t} + q_{2xx} + 2(|q_1|^2 + |q_2|^2)q_2 = 0; \quad (1b)$$

The integrability of this system was proved by Manakov, which is here named as the integrable Manakov^[6] system. Recently, the rogue wave phenomenon was popular in nonlinear science. Rogue waves are strong wavelets that may appear in the ocean when special conditions are met. These waves can be two, three or even more times higher than the average wave crest. Rogue waves are also known as freak waves, monster waves, killer waves, giant waves or extreme waves.^[10] The first-order rational solution for the nonlinear Schrödinger equation was given by Peregrine in 1983.^[11] The rogue wave of the NLS equation was derived by Darboux transformation.^[12] Recently, we constructed some general rogue wave solutions by the Darboux-Liouville transformation.^[13] It is well known that there have been lots of results about the rogue wave for nonlinear Schrödinger equations. However, to the best of our knowledge, there are no reports on exact solutions related to the rogue wave solution of the coupled Schrödinger equation. Here we also obtain the new breather solutions and bright-dark-rogue solutions as a by-product.

The Darboux transformation^[14–17] is a powerful method to construct some interesting solutions (solitons, positons, rogue wave solution) in the integrable system.

The goal of this work is to use the Darboux transformation to construct the rogue wave solution, breathers and bright-dark-rogue solution. These special solutions may play an important role in the research of some physical phenomena such as BEC or nonlinear optics.

Firstly, we construct the rogue wave solution by the Darboux transformation. The system (1) admits the following Lax pair

$$\Phi_x = U\Phi = [i \ U_1 + U_0]\Phi; \quad (2a)$$

$$\Phi_t = V\Phi = [3i \ U_1 + 3 \ U_0 + i \ U_3(U_{0,x} - U_0^2)]\Phi; \quad (2b)$$

where

$$U_0 = \begin{pmatrix} 0 & q_1 & q_2 \\ -q_1^* & 0 & 0 \\ -q_2^* & 0 & 0 \end{pmatrix};$$

$U_1 = \text{diag}\{-2; 1; 1\}$; $U_3 = \text{diag}\{1; -1; -1\}$; and Φ is a 3×3 -matrix valued eigenfunction, λ is a spectral parameter. The compatibility condition of the Lax pair (2) gives the CNLS (1). It is well known that the Darboux transformation^[7,15,16] for this system (2) is

$$T = I - [\lambda^* - (\lambda^* - 1)P_1]; \quad P_1 = \frac{\Phi_1 \Phi_1^\dagger}{\Phi_1^\dagger \Phi_1}; \quad (3)$$

where $\Phi_1 = \Phi(x; t; \lambda_1)(m_1; m_2; m_3)^T$, $m_1; m_2; m_3$ are constants, I is a 3×3 identity matrix and $\Phi(x; t; \lambda_1)$ is the fundamental solution for the Lax pair equation at $\lambda = \lambda_1$, $q_i = q_i[0]$ ($i = 1, 2$). It follows the transformation between the fields,

$$\begin{aligned} q_1[1] &= q_1[0] + 6\text{Im}(\lambda_1) \frac{1}{|q_1|^2 + |q_2|^2 + |q_3|^2}; \\ q_2[1] &= q_2[0] + 6\text{Im}(\lambda_1) \frac{1}{|q_1|^2 + |q_2|^2 + |q_3|^2}; \end{aligned} \quad (4)$$

where $(\lambda_1; \lambda_2; \lambda_3)^T = \Phi_1$.

**Email: lingliming@qq.com

© 2011 Chinese Physical Society and IOP Publishing Ltd

We start from the seed solution-nonzero plane wave solution. We denote them as

$$q_1[0] = c_1 \exp[i \varphi_1]; \quad q_2[0] = c_2 \exp[i \varphi_2]; \quad (5)$$

where $\varphi_i = d_i x + (2c_1^2 + 2c_2^2 - d_i^2)t$ with c_i and d_i ($i = 1, 2$) being arbitrary constants. However, these Lax pair equations are variable coefficient differential equations. We can not directly solve them. Luckily we can convert this system into constant coefficient differential equations by a gauge transformation. In the following, we give the transformation for the Lax pair equations (2) at $q_1 = q_1[0]$ and $q_2 = q_2[0]$. We can obtain

$$\begin{aligned} \Psi_x &= (MUM^{-1} + M_x M^{-1})\Psi = U_1 \Psi; \\ \Psi_t &= (MV M^{-1} + M_t M^{-1})\Psi = V_1 \Psi; \end{aligned} \quad (6)$$

where $\Psi = M\Phi$, $M = \text{diag}\{\exp[-\frac{i}{3}(\varphi_1 + \varphi_2)]; \exp[\frac{i}{3}(2\varphi_1 - \varphi_2)]; \exp[\frac{i}{3}(2\varphi_2 - \varphi_1)]\}$. Simply computing, we have

$$U_1 = \begin{pmatrix} 11 & c_1 & c_2 \\ -c_1 & 22 & 0 \\ -c_2 & 0 & 33 \end{pmatrix};$$

$$V_1 = iU_1^2 - [\frac{2}{3}(d_1 + d_2) - 2\varphi_1]U_1 + mI; \quad (7)$$

$$\begin{aligned} 11 &= -2i\varphi_1 - \frac{i}{3}(d_1 + d_2); \\ 22 &= i\varphi_1 + \frac{i}{3}(2d_1 - d_2); \\ 33 &= i\varphi_1 + \frac{i}{3}(2d_2 - d_1); \end{aligned}$$

where $m = 2i\varphi_1 + \frac{2i}{3}(c_1^2 + c_2^2) + \frac{2i}{9}(d_1^2 - d_1 d_2 + d_2^2) + \frac{2i\varphi_1}{3}(d_1 + d_2)$. In Refs. [7–9], the authors consider the solution when the characteristic equation for U_1 has a single and double root. Indeed, the characteristic equation for U_1 can possess a triple root. Our work is looking for the solution for this special assumption. Equations (6) are the constant coefficient differential equations. In order to look for the “rational solution”, we choose the parameters to satisfy $\varphi_1 = d_2 + 3e_1$; $e_1 = \text{Re}(\varphi_1)$; $d_1 = d_2 - 2$; $c_1 = \pm 2$; $c_2 = \pm 2$; $\text{Im}(\varphi_1) = \pm\sqrt{3}$, d_2 and $\text{Re}(\varphi_1)$ are arbitrary real numbers. The matrix U_1 cannot diagonalize and can merely decompose with the Jordan form with the diagonal element. Therefore the fundamental-solution matrix for Lax pair equations (2) at $\varphi_i = \varphi_i[0]$, $q_i = q_i[0]$ ($i = 1, 2$) are $\Phi(x; t; \varphi_i) = M^{-1}\Theta$;

$$\Theta = \begin{pmatrix} \$_{11} & 4\varphi_1^2 + 2\sqrt{3} & 4 \\ \$_{12} & -2\varphi_1^2(\sqrt{3} - i) & -2 & -2\varphi_1^2(\sqrt{3} - i) \\ \$_{13} & -2\varphi_1^2(\sqrt{3} + i) & -2 & -2\varphi_1^2(\sqrt{3} + i) \end{pmatrix};$$

$$\begin{aligned} \$_{11} &= 4\varphi_1^2(\varphi_1^2 + 2it) + 4\sqrt{3} & -2; \\ \$_{12} &= -2(\sqrt{3} - i)\varphi_1^2(\varphi_1^2 + 2it) - 4 & ; \\ \$_{13} &= -2(\sqrt{3} + i)\varphi_1^2(\varphi_1^2 + 2it) - 4 & ; \end{aligned}$$

where $\varphi_1 = x + 2\sqrt{3}it$; and $\varphi_2 = -i\sqrt{3}e_1$. In what follows, we use the Darboux transformation (4) to derive the rational solution for CNLS (1).

(1) *Rogue wave I*. Choosing the parameters $m_1 = 0$, $m_2 = 1$, $m_3 = 0$, we obtain the one rogue wave I by the formula (4)

$$\begin{aligned} q_1[1] &= \left[-1 - i\sqrt{3} \right. \\ &\quad \left. + \frac{-6\sqrt{3} - 36t^2\sqrt{3} - 3 + i(36\varphi_1^2 t + 6\varphi_1^2 + 5\sqrt{3})}{12\varphi_1^2 t^2 + 8\sqrt{3} + 144t^4 + 5} \right] \\ &\quad \cdot \exp[i\varphi_1]; \\ q_2[1] &= \left[-1 + i\sqrt{3} \right. \\ &\quad \left. + \frac{-6\sqrt{3} + 36t^2\sqrt{3} - 3 + i(36\varphi_1^2 t - 6\varphi_1^2 - 5\sqrt{3})}{12\varphi_1^2 t^2 + 8\sqrt{3} + 144t^4 + 5} \right] \\ &\quad \cdot \exp[i\varphi_2]; \end{aligned} \quad (8)$$

where $\varphi_1 = x + 6e_1 t$. This kind of rogue wave solution is similar to the first-order rogue wave solution for the nonlinear Schrödinger equation.^[12]

(2) *Rogue wave II*. Choosing the parameters $m_1 = 1$, $m_2 = 0$, $m_3 = 0$, we obtain the rogue wave II by the formula

$$\begin{aligned} q_1[1] &= [-1 - i\sqrt{3} + \frac{G_1 + iH_1}{D}] \exp[i\varphi_1]; \\ q_2[1] &= [-1 + i\sqrt{3} + \frac{G_2 + iH_2}{D}] \exp[i\varphi_2]; \end{aligned} \quad (9)$$

where

$$\begin{aligned} D &= 1728\varphi_1^8 t^4 + 288\varphi_1^6 t^2 + 384\sqrt{3}\varphi_1^5 t^2 + 12\varphi_1^4 t^4 \\ &\quad + 432\varphi_1^4 t^2 + 16\sqrt{3}\varphi_1^3 t^3 + 24\varphi_1^2 t^2 + 4\sqrt{3}\varphi_1 + 1; \\ G_1 &= -864\sqrt{3}\varphi_1^6 t^3 - 144\sqrt{3}\varphi_1^5 t^2 - 72\sqrt{3}\varphi_1^4 t^2 t \\ &\quad - 216\varphi_1^4 t^2 - 12\sqrt{3}\varphi_1^3 t^3 - 144\varphi_1^3 t - 18\varphi_1^2 t^2 \\ &\quad - 12\sqrt{3}\varphi_1^2 t + 3; \\ G_2 &= +864\sqrt{3}\varphi_1^6 t^3 - 144\sqrt{3}\varphi_1^5 t^2 + 72\sqrt{3}\varphi_1^4 t^2 t \\ &\quad - 216\varphi_1^4 t^2 - 12\sqrt{3}\varphi_1^3 t^3 + 144\varphi_1^3 t - 18\varphi_1^2 t^2 \\ &\quad + 12\sqrt{3}\varphi_1^2 t + 3; \\ H_1 &= +864\varphi_1^6 t^3 + 144\varphi_1^5 t^2 + 72\varphi_1^4 t^2 t + 312\sqrt{3}\varphi_1^4 t^2 \\ &\quad + 12\varphi_1^3 t^3 + 96\sqrt{3}\varphi_1^3 t + 18\sqrt{3}\varphi_1^2 t^2 + 108\varphi_1^2 t \\ &\quad + 12\varphi_1 + \sqrt{3}; \\ H_2 &= +864\varphi_1^6 t^3 - 144\varphi_1^5 t^2 + 72\varphi_1^4 t^2 t - 312\sqrt{3}\varphi_1^4 t^2 \\ &\quad - 12\varphi_1^3 t^3 + 96\sqrt{3}\varphi_1^3 t - 18\sqrt{3}\varphi_1^2 t^2 + 108\varphi_1^2 t \\ &\quad - 12\varphi_1 - \sqrt{3}. \end{aligned}$$

Choosing the right parameter, the pictures for $|q_1[1]|^2$ and $|q_2[2]|^2$ are shown in Fig. 1. They have different behaviors to rogue waves of NLS.^[12] Indeed we can combine the three vector solution to obtain the new rogue wave, but we cannot obtain any new kinds of

solution except the above two kinds. The rogue wave we obtained is different from Ref. [4] by a symmetry analysis, which is nothing but the rogue wave of NLS at different altitude.




Fig. 1. Different behavior of $|q_1|^2$ and $|q_2|^2$ besides the altitude for the parameters $c_1 = \frac{1}{10}$, $d_1 = 0$.




Fig. 2. (a) and (b) with parameters $c = -2$, $d = 0$, $\zeta_1 = 2i$, $\lambda_1 = 1 - 5$, $\tau_1 = \tau_2 = \kappa_1 = \kappa_2 = 0$; and the bright-dark-rogue solution [(c) and (d)] with parameters $c = d = 1$, $c_1 = c_2 = c_3 = 1$.

We can easily obtain the following relation for those rogue wave solutions,

$$\begin{aligned} \int_{-\infty}^{\infty} (|q_1[1]|^2 - 4^{-2}) dx &= 0; \\ \int_{-\infty}^{\infty} (|q_2[1]|^2 - 4^{-2}) dx &= 0; \end{aligned} \quad (10)$$

This means that $|q_1[1]|^2$ and $|q_2[1]|^2$ are equally above and below the background.

Moreover, we consider another interesting case: one is a zero solution and the other is a plane wave solution. It is interesting that we could obtain breather solutions, a bright-dark solution and a bright-dark-rogue solution by this choice. In what follows, we start from the seed solution $q_1[0] = c \exp[i \cdot]$, $q_2[0] = 0$; where $\cdot = dx + (2c^2 - d^2)t$. Similarly, we take

$M = \text{diag}\{1; \exp[i \cdot]; 1\}$. Then we can obtain the following results

$$U_1 = \begin{pmatrix} -2i & 1 & c & 0 \\ -c & i & 1 + id & 0 \\ 0 & 0 & 0 & i & 1 \end{pmatrix};$$

$$V_1 = iU_1^2 + 2 \cdot U_1 + 2i(\cdot^2 + c^2) - \text{diag}\{0; 0; 2ic^2\};$$

The characteristic equation for U_1 is $(-i \cdot)(\cdot + 2i \cdot)(-i \cdot - id) + c^2 = 0$. Then we would classify the characteristic equation into two kinds that different solution behaviors will exhibit.

Case I: single root. Assume that the characteristic equation of matrix U_1 has three distinct roots. Denote them as $i \cdot$, $\cdot = \frac{i}{2}[-\cdot + d + \sqrt{9 \cdot^2 + 6 \cdot d + d^2 + 4c^2}]$, $\cdot = i(d - \cdot) - \cdot$. Finally, we obtain the fundamental solution for the Lax pair equations $\Phi(x; t; \cdot) = M^{-1}Q\Omega$;

$$Q = \begin{pmatrix} 0 & c & \cdot + 2i \cdot & 1 \\ 0 & \cdot + 2i \cdot & c & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix};$$

$$\Omega = \begin{pmatrix} e^{i\zeta_1 x + 3i\zeta_1^2 t} & 0 & 0 \\ 0 & 22 & 0 \\ 0 & 0 & 33 \end{pmatrix};$$

$$22 = \exp\{\cdot x + [i \cdot^2 + 2 \cdot \cdot + 2i(\cdot^2 + c^2)]t\};$$

$$33 = \exp\{\cdot x + [i \cdot^2 + 2 \cdot \cdot + 2i(\cdot^2 + c^2)]t\};$$

Substituting the solution to the formula (4) and taking the parameters $m_1 = 1$, $m_2 = e^{\tau_1 + i\kappa_1}$, $m_3 = e^{\tau_2 + i\kappa_2}$, ($\cdot$, $\cdot$ ($i = 1; 2$) are real numbers), we could obtain the breather solutions for the CNLS equation. This kind of solution is a periodic function or a quasi-periodic function. Taking the proper parameters, then Figs. 2(a) and 2(b) exhibit the dynamical behavior of the solutions $q_1[1]$ and $q_2[1]$. When $\text{Re}(\cdot)$

it follows that $q_1[1]$ is the rogue wave solution for the NLS equation and $q_2[1] = 0$. In the following, we consider the other two non-trivial cases.

Case 1: Taking the parameters $m_1 = 1$, $m_2 = 0$, $m_3 = a_3 + b_3i$, we obtain the dark-bright solitons,

$$q_1[1] = c \tanh\left(\frac{3}{2}x_1 + \dots\right) \exp(i \dots);$$

$$q_2[1] = -4c^2(a_3 + ib_3) \operatorname{sech}\left(\frac{3}{2}x_1 + \dots\right) \cdot \exp[i(dx + (3c^2 - d^2)t)];$$

where $x_1 = \frac{2c}{3}(x - 2dt)$ and $\dots = \frac{1}{2} \ln\left(\frac{a_3^2 + b_3^2}{2c^2}\right)$. It is interesting that we can obtain the bright-dark soliton solution in the focusing behavior and anomalous dispersion fibers.

Case 2: Taking parameters $m_1 = a_1 + b_1i$, $m_2 = 1$, $m_3 = a_3 + b_3i$, we have

$$q_1[1] = c \left\{ 1 - 4 \left[\frac{x_1^2}{2} + \frac{x_2^2}{2} - x_1 + i x_2 \right] \exp(-x_1) \cdot \left\{ 2 \left(\frac{x_1^2}{2} + \frac{x_2^2}{2} \right) - 2x_1 + 1 \right\} \exp(-x_1) + (a_3^2 + b_3^2) \exp(2x_1) \right\}^{-1} \exp(i \dots);$$

$$q_2[1] = -4c \left[a_3(x_1 - 1) - b_3 x_2 + i[a_3 x_2 + b_3(x_1 - 1)] \right] \cdot \exp\left[\frac{1}{2}x_1 + i(dx + (3c^2 - d^2)t)\right] \cdot \left\{ 2 \left(\frac{x_1^2}{2} + \frac{x_2^2}{2} \right) - 2x_1 + 1 \right\} \exp(-x_1) + (a_3^2 + b_3^2) \exp(2x_1) \right\}^{-1};$$

where $x_1 = a_1c + c(x - 2dt)$, $x_2 = b_1c - 2c^2t$. We plot the picture for this kind of solution by choosing the

right parameters (Figs. 2(c) and 2(d)). This kind of solution behaves like the nonlinear superposition for the rogue wave solution and bright-dark soliton solution.

References

- [1] Agrawal G P 1995 *Nonlinear Fiber Optics* (New York: Academic)
- [2] Scott A C 1984 *Physica Scripta* **29** 279
- [3] E P Bashkin and Vagov A V 1997 *Phys. Rev. B* **56** 6207
- [4] Yan Z 2011 *Financial Rogue Waves Appearing in the Coupled Nonlinear Volatility and Option Pricing Model* arxiv:1101.3107v1
- [5] K Dysthe, Krogstad H E and Müller P 2008 *Annu. Rev. Fluid Mech.* **40** 287
- [6] Manakov S V 1973 *ZETP* **65** 505
- [7] Forest M G, McLaughlin D W, Muraki D J and Wright O C 2000 *J. Non. Sci.* **10** 291
- [8] Wright O C and Forest M G 2000 *Physica D* **141** 104
- [9] Forest M G, Sheu S P and Wright O C 2000 *Phys. Lett. A* **266** 24
- [10] Osborne A R 2009 *Nonlinear Ocean Waves* (New York: Academic)
- [11] Peregrine D H 1983 *J. Sust. Math. Soc. B: Appl. Math.* **25** 16
- [12] Akhmediev N, Ankiewicz A and Soto-Crespo J M 2009 *Phys. Rev. E* **80** 026601
- [13] Guo B and Ling L 2011 *Acta Mathematica Scientia* (submitted)
- [14] Matveev V B and Salle M A 1991 *Darboux Transformations and Solitons* (New York: Springer)
- [15] Gu C H, Hu H S and Zhou Z X 2005 *Darboux Transformations in Integrable Systems: Theory and Their Applications* (Berlin: Springer)
- [16] Doktorov E V and Leble S B 2007 *A Dressing Method in Mathematical Physics* (New York: Springer)
- [17] Ling L and Liu Q P 2010 *J. Phys. A: Math. Theor.* **43** 434023

Chinese Physics Letters

Volume 28

Number 11

November 2011

GENERAL

- 110201 **Inertial Levy Flight with Nonlinear Friction**
LÜ Yan, BAO Jing-Dong
- 110202 **Rogue Wave, Breathers and Bright-Dark-Rogue Solutions for the Coupled Schrödinger Equations**
GUO Bo-Ling, LING Li-Ming
- 110203 **Conformal Invariance of Higher-Order Lagrange Systems by Lie Point Transformation**
HUANG Wei-Li, CAI Jian-Le
- 110204 **Analysis and Control of Two-Layer Frenkel-Kontorova Model**
TANG Wen-Yan, QU Zhi-Hua, GUO Yi
- 110205 **Novel Approach for a van der Pol Oscillator in the Continuous Time Domain**
Junaid Ali Khan, Muhammad Asif Zahoor Raja, Ijaz Mansoor Qureshi
- 110301 **Dynamics of Nonclassical Correlation in Interacting Qubits under Correlated Dissipative Environments**
QIU Liang
- 110401 **Unification of General Relativity with Quantum Field Theory**
NI Jun
- 110501 **Random Long-Range Interaction Induced Synchronization in Coupled Networks of Inertial Ratchets**
WEI Du-Qu, LUO Xiao-Shu, CHEN Hong-Bin, ZHANG Bo
- 110502 **Effect of Interface Structure on Thermal Boundary Conductance by using First-principles Density Functional Perturbation Theory**
GAO Xue, ZHANG Yue, SHANG Jia-Xiang
- 110503 **Fractal Erosion of the Safe Basin in a Helmholtz Oscillator and Its Control by Linear Delayed Velocity Feedback**
SHANG Hui-Lin, WEN Yong-Peng
- 110504 **Predicting Natural and Chaotic Time Series with a Swarm-Optimized Neural Network**
Juan A. Lazzús
- 110505 **Vibrational and Stochastic Resonance in the FitzHugh-Nagumo Neural Model with Multiplicative and Additive Noise**
HE Zheng-You, ZHOU Yu-Rong
- 110506 **Chaos Suppression in a Sine Square Map through Nonlinear Coupling**
Eduardo L. Brugnago, Paulo C. Rech
- 110507 **Adaptive Increasing-Order Synchronization and Anti-Synchronization of Chaotic Systems with Uncertain Parameters**
M. Mossa Al-sawalha, M. S. M. Noorani
- 110601 **Temporal Parameter Optimization in Four-Dimensional Flash Trajectory Imaging**
WANG Xin-Wei, ZHOU Yan, FAN Song-Tao, LIU Yu-Liang
- 110701 **Highly Sensitive Rayleigh Wave Hydrogen Sensors with WO_3 Sensing Layers at Room Temperature**
WANG Cheng, FAN Li, ZHANG Shu-Yi, YANG Yue-Tao, ZHOU Ding-Mao, SHUI Xiu-Ji
- 110702 **Pressure-induced Phase Transition in Oleic Acid Studied by Raman Spectroscopy**
FAN Ya, ZHOU Jing, LI Shuang, GUAN Fu-Ying, XU Da-Peng

THE PHYSICS OF ELEMENTARY PARTICLES AND FIELDS

**111201 Fermion Mass Hierarchies in Singlet-Extended
Minimal-Supersymmetric-Standard-Model Quivers**

Salah Eddine Ennadifi, Adil Belhaj, El Hassan Saidi

111202

JUST FOR AUTHORS
— CHINESE PHYSICS LETTERS

- 114207 An InP-Based Dual-Depletion-Region Electroabsorption Modulator with Low Capacitance and Predicted High Bandwidth**
SHAO Yong-Bo, ZHAO Ling-Juan, YU Hong-Yan, QIU Ji-Fang, QIU Ying-Ping, PAN Jiao-Qing, WANG Bao-Jun, ZHU Hong-Liang, WANG Wei
- 114208 Diode Laser Optically Injected by Resonance of a Monolithic Cavity**
PENG Yu, ZHAO Yang, LI Ye, YANG Tao, CAO Jian-Ping, FANG Zhan-Jun, ZANG Er-Jun
- 114209 Automatic Compensation of Total Phase Aberrations in Digital Holographic Biological Imaging**
ZHANG Yi-Zhuo, WANG Da-Yong, WANG Yun-Xin, TAO Shi-Quan
- 114210 Novel Propagation Properties of Total Internal Reflection Photonic Crystal Fibres with Rhombic Air Holes**
YAN Hai-Feng, YU Zhong-Yuan, LIU Yu-Min, TIAN Hong-Da, HAN Li-Hong
- 114211 Single and Multicasting Inverted-Wavelength Conversion at 80 Gb/s Based on a Single Semiconductor Optical Amplifier**
HUANG Xi, QIN Cui, YU Yu, ZHANG Xin-Liang
- 114212 Multicolor InAs/InP(100) Quantum Dot Laser**
LI Shi-Guo, GONG Qian, CAO Chun-Fang, WANG Xin-Zhong, WANG Rui-Chun, YUE Li, LIU Qing-Bo, WANG Hai-Long
- 114301 Resonance Effects of Bilayered Piezoelectric Films Used for Bulk Acoustic Wave Sensors**
ZHANG Hui, ZHANG Shu-Yi, FAN Li
- 114302 Data-Derived Estimation of Source Depth Using Vertical Line Array Data in Shallow Water**
WANG Hao-Zhong, WANG Ning, GAO Da-Zhi
- 114303 Analytical Solution of the Blast Wave Problem in a Non-Ideal Gas**
L. P. Singh, S. D. Ram, D. B. Singh
- 114701 Effects of a Premixed Layer on the Richtmyer-Meshkov Instability**
TIAN Bao-Lin, ZHANG Xin-Ting, QI Jin, WANG Shuang-Hu
- 114702 Viscous Swirling Flow over a Stretching Cylinder**
Tiegang FANG, Shanshan YAO

PHYSICS OF GASES, PLASMAS, AND ELECTRIC DISCHARGES

- 115201 Observation of Hot Electrons in Surface-Wave Plasmas Excited by Surface Plasmon Polaritons**
HU Ye-Lin, CHEN Zhao-Quan, LIU Ming-Hai, HONG Ling-Li, LI Ping, ZHENG Xiao-Liang, XIA Guang-Qing, HU Xi-Wei
- 115202 Electric Discharge in Pin-Plate Audio Frequency Plasma**
A. A. Azooz, S. K. Talal

CONDENSED MATTER: STRUCTURE, MECHANICAL AND THERMAL PROPERTIES

- 116101 Optical Limiting Properties of Ag-Cu Metal Alloy Nanoparticles Analysis by using MATLAB**
WANG Yu-Hua, LI Hui-Qing, LU Jian-Duo, WANG Ru-Wu
- 116102 Formation Mechanism of Micropores on the Surface of Pure Aluminum Induced by High-Current Pulsed Electron Beam Irradiation**
ZOU Yang, CAI Jie, WAN Ming-Zhen, LV Peng, GUAN Qing-Feng
- 116103 Structural and Magnetic Studies of $\text{Zn}_{0.95}\text{Co}_{0.05}\text{O}$ and $\text{Zn}_{0.90}\text{Co}_{0.05}\text{Al}_{0.05}\text{O}$**
Murtaza Saleem, Saadat A. Siddiqi, Shahid Atiq, M. Sabieh Anwar
- 116104 Effects of the Cooling Rate on the Plasticity of $\text{Pd}_{40.5}\text{Ni}_{40.5}\text{P}_{19}$ Bulk Metallic Glasses**
LI Yang, QIU Sheng-Bao, SHAO Yang, YAO Ke-Fu

- 116201 Stiction of a Nano-Beam with Surface Effect
LIU Jian-Lin, XIA Re, ZHOU Yue-Ting
- 116801 Hydrogen Storage Materials Based on Single-Layer Aluminum Nitride Nanostructures
WANG Yu-Sheng, YUAN Peng-Fei, LI Meng, SUN Qiang, JIA Yu
- 116802 Effect of Interface Bond Type on the Structure of InAs/GaSb Superlattices Grown by Metalorganic Chemical Vapor Deposition
LI Li-Gong, LIU Shu-Man, LUO Shuai, YANG Tao, WANG Li-Jun, LIU Feng-Qi, YE Xiao-Ling, XU Bo, WANG Zhan-Guo
- 116803 Simulation and Suppression of the Gas Phase Pre-reaction in Metal-Organic Chemical Vapor Deposition of ZnO
ZHU Guang-Yao, GU Shu-Lin, ZHU Shun-Ming, TANG Kun, YE Jian-Dong, ZHANG Rong, SHI Yi, ZHENG You-Dou
- 116804

JUST FOR AUTHORS
— CHINESE PHYSICS LETTERS

- 118102 **Annealing Effect of Pulsed Laser Deposited Transparent Conductive Ta-Doped Titanium Oxide Films**
WU Bin-Bin, PAN Feng-Ming, YANG Yu-E
- 118103 **Mechanical Performances of Carbonitriding Films on Cast Iron by Plasma Electrolytic Carbonitriding**
PANG Hua, ZHANG Gu-Ling, WANG Xing-Quan, LV Guo-Hua, CHEN Huan, YANG Si-Ze
- 118104 **Preparation and Luminescence Properties of the $\text{Na}_2\text{W}_2\text{O}_7$ Crystal**
WEI Qin-Hua, SHI Hong-Sheng, CHAI Wen-Xiang, QIN Lai-Shun, SHU Kang-Ying
- 118105 **Efficiency Droop Effect Mechanism in an InGaN/GaN Blue MQW LED**
WANG Jia-Xing, WANG Lai, HAO Zhi-Biao, LUO Yi
- 118301 **Bifurcation of a Swelling Gel with a Mechanical Load and Geometric Constraint**
XUE Feng, YONG Hua-Dong, ZHOU You-He
- 118401 **Partially Loaded Cavity Analysis by Using the 2-D FDTD Method**
YAO Bin, ZHENG Qin-Hong, PENG Jin-Hui, ZHONG Ru-Neng, XIANG Tai, XU Wan-Song
- 118501 **Improved Performance of Fluorinated Copper Phthalocyanine Thin Film Transistors Using Para-hexaphenyl as the Inducing Layer**
MA Feng, WANG Shi-Rong, LI Xiang-Gao, YAN Dong-Hang
- 118502 **High Performance Polymer Field-Effect Transistors Based on Thermally Crosslinked Poly(3-hexylthiophene)**
JIANG Chun-Xia, YANG Xiao-Yan, ZHAO Kai, WU Xiao-Ming, YANG Li-Ying, CHENG Xiao-Man, WEI Jun, YIN Shou-Gen
- 118701 **Depth Determination of an Abnormal Heat Source in Biological Tissues**
WANG Qing-Hua, LI Zhen-Hua, LAI Jian-Cheng, HE An-Zhi
- 118702 **Effects of Labeling Thiophilic FRET Dyes on the Stability and Dimerization Process of α -Lactoglobulin**
PAN Hai, XIE Jin-Bing, CAO Yi, QIN Meng, WANG Wei
- 118901 **Self-Organized Criticality in Small-World Networks Based on the Social Balance Dynamics**
MENG Qing-Kuan
- 118902 **Percolation of Mobile Individuals on Weighted Scale-Free Networks**
WU An-Cai
- 118903 **Interactions between Intersecting Pedestrian and Vehicle Flows on Roads**
GUO Ren-Yong, GUO Xi

GEOPHYSICS, ASTRONOMY, AND ASTROPHYSICS

- 119601 **Model of the Crustal Magnetic Field in the Martian Aurora Zone**
TONG Dong-Sheng, CHEN Chu-Xin
- 119801 **Modified Chaplygin Gas with Variable G and Λ**
Ujjal Debnath

JUST FOR AUTHORS
— CHINESE PHYSICS LETTERS